

84°41'43"W



84° 41' 43" W

600

Map projection: Web Mercator Corner coordinates: WGS84

1200

100

300

0

 \mathbb{N}

USDA

Web Soil Survey National Cooperative Soil Survey

____Feet 1800

4/10/2014 Page 1 of 5

		MA	AP LEGEND			MAP INFORMATION
Area of Int	erest (AOI) Area of Interest (AOI)	~	Loamy, siliceous, subactive, thermic Arenic Plinthaquic Paleudults	%	Major Roads Local Roads	The soil surveys that comprise your AOI were mapped at 1:12,000.
Soils Soil Pat	ng Polygons	~	Sandy, siliceous, thermic Cumulic Humaguepts	Backgrou		Warning: Soil Map may not be valid at this scale.
	Fine, kaolinitic, thermic		Not rated or not available	ALC: NO	Aerial Photography	Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil
	Typic Paleaquults	Soil Rat	ing Points			line placement. The maps do not show the small areas of
	Fine-loamy, kaolinitic, thermic Plinthic Kandiudults		Fine, kaolinitic, thermic Typic Paleaquults			contrasting soils that could have been shown at a more detailed scale.
	Fine-loamy, kaolinitic, thermic Typic Kandiudults		Fine-loamy, kaolinitic, thermic Plinthic Kandiudults			Please rely on the bar scale on each map sheet for map measurements.
	Loamy, siliceous, subactive, thermic Arenic Plinthaquic Paleudults		Fine-loamy, kaolinitic, thermic Typic Kandiudults			Source of Map: Natural Resources Conservation Service
	Sandy, siliceous, thermic Cumulic Humaquepts		Loamy, siliceous, subactive, thermic Arenic Plinthaguic Paleudults			Web Soil Survey URL: http://websoilsurvey.nrcs.usda.gov Coordinate System: Web Mercator (EPSG:3857)
Soil Rat	Not rated or not available ng Lines		Sandy, siliceous, thermic Cumulic Humaquepts			Maps from the Web Soil Survey are based on the Web Mercato projection, which preserves direction and shape but distorts
~	Fine, kaolinitic, thermic Typic Paleaquults		Not rated or not available			distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.
~	Fine-loamy, kaolinitic,	Water Fea	Itures Streams and Canals			
	thermic Plinthic Kandiudults	Transport				This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.
-	Fine-loamy, kaolinitic,	+++	Rails			Soil Survey Area: Gadsden County, Florida
	thermic Typic Kandiudults	~	Interstate Highways			Survey Area Data: Version 19, Dec 27, 2013
		~	US Routes			Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.
						Date(s) aerial images were photographed: Apr 4, 2010—Jan 12, 2011
						The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

#3 Farm

Soil Taxonomy Classification

100.0%	159.0		st	Totals for Area of Interest
1.5%	2.4	Loamy, siliceous, subactive, thermic Arenic Plinthaquic Paleudults	Leefield fine sand, 0 to 5 percent slopes	113
9.5%	15.2	Sandy, siliceous, thermic Cumulic Humaquepts	Pickney, Dorovan, and Bibb soils, frequently flooded	66
41.9%	66.7	Fine-loamy, kaolinitic, thermic Typic Kandiudults	Fuquay-Orangeburg- Norfolk complex, 8 to 15 percent slopes	48
18.7%	29.7	Fine-loamy, kaolinitic, thermic Typic Kandiudults	Orangeburg loamy sand, 2 to 5 percent slopes	46
3.0%	4.7	Fine-loamy, kaolinitic, thermic Typic Kandiudults	Orangeburg loamy sand, 0 to 2 percent slopes	45
4.5%	7.1	Loamy, siliceous, subactive, thermic Arenic Plinthaquic Paleudults	Leefield-Bonifay-Dothan complex, 0 to 5 percent slopes	32
2.6%	4.1	Fine, kaolinitic, thermic Typic Paleaquults	Grady fine sandy loam, depressional	29
14.6%	23.1	Fine-loamy, kaolinitic, thermic Plinthic Kandiudults	Dothan-Fuquay complex, 2 to 5 percent slopes	19
3.7%	5.9	Fine-loamy, kaolinitic, thermic Plinthic Kandiudults	Dothan-Fuquay complex, 0 to 2 percent slopes	18
Percent of AOI	Acres in AOI	Rating	Map unit name	Map unit symbol
(FL039)	Gadsden County, Florida	Summary by Map Unit —	Soil Taxonomy Classification— Summary by Map Unit — Gadsden County, Florida (FL039)	Soil Tay

USDA

#3 Farm

Description

This rating presents the taxonomic classification based on Soil Taxonomy

Classification is based on soil properties observed in the field or inferred from those observations or from laboratory measurements. This table shows the classification The system of soil classification used by the National Cooperative Soil Survey has six categories (Soil Survey Staff, 1999 and 2003). Beginning with the broadest, paragraphs. of the soils in the survey area. The categories are defined in the following these categories are the order, suborder, great group, subgroup, family, and series

is identified by a word ending in sol. An example is Alfisols. the dominant soil-forming processes and the degree of soil formation. Each order ORDER. Twelve soil orders are recognized. The differences among orders reflect

name of a suborder indicates the order. An example is Udalfs (Ud, meaning humid plus alfs, from Alfisols). that reflect the most important variables within the orders. The last syllable in the properties that influence soil genesis and are important to plant growth or properties SUBORDER. Each order is divided into suborders primarily on the basis of

regime). horizonation, plus udalfs, the suborder of the Alfisols that has a udic moisture indicates a property of the soil. An example is Hapludalfs (Hapl, meaning minimal status. Each great group is identified by the name of a suborder and by a prefix that horizons; soil moisture and temperature regimes; type of saturation; and base similarities in kind, arrangement, and degree of development of pedogenic GREAT GROUP. Each suborder is divided into great groups on the basis of close

orders, suborders, or great groups. Extragrades have some properties that are not group; it is not necessarily the most extensive. Intergrades are transitions to other the great group. An example is Typic Hapludalfs. the name of the great group. The adjective Typic identifies the subgroup that typifies taxonomic class. Each subgroup is identified by one or more adjectives preceding representative of the great group but do not indicate transitions to any other intergrades or extragrades. The typic subgroup is the central concept of the great SUBGROUP. Each great group has a typic subgroup. Other subgroups are

fine-loamy, mixed, active, mesic Typic Hapludalfs. name of a subgroup preceded by terms that indicate soil properties. An example is temperature regime, soil depth, and reaction class. A family name consists of the particle-size class, mineralogy class, cation-exchange activity class, soil biological activity. Among the properties and characteristics considered are the properties are those of horizons below plow depth where there is much chemical properties and other characteristics that affect management. Generally, FAMILY. Families are established within a subgroup on the basis of physical and

and arrangement in the profile. color, texture, structure, reaction, consistence, mineral and chemical composition, SERIES. The series consists of soils within a family that have horizons similar in

USDA

References:

Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service. U.S. Department of Agriculture Handbook 436.

Soil Survey Staff. 2006. Keys to soil taxonomy. 10th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. (The soils in a given survey area may have been classified according to earlier editions of this publication.)

Rating Options

Aggregation Method: Dominant Condition Component Percent Cutoff: None Specified Tie-break Rule: Lower