Soils Map



Weighted Average						148.9	43.2	54.7	*-	63.5	63.4	29.8
980B	Gullied land-Ely-Colo complex, 2 to 5 percent slopes	0.08	0.2%		VIIe	88	25.5	42	25	32	5	0
Y93D2	Shelby-Adair clay loams, dissected till plain, 9 to 14 percent slopes, eroded	0.69	1.5%		llle	0	0	41		60	60	37
370C2	Sharpsburg silty clay loam, 5 to 9 percent slopes, eroded	1.32	2.9%		llle	204.8	59.4	80	67	71	71	48
Y428B	Ely silty clay loam, dissected till plain, 2 to 5 percent slopes	2.34	5.1%		lle	0	0	88		96	96	31
822D3	Lamoni soils, 9 to 14 percent slopes, severely eroded	4.57	9.9%		Vle	89.6	26	5	5	36	36	12
54	Zook silty clay loam, 0 to 2 percent slopes, occasionally flooded	4.69	10.1%		llw	164.8	47.8	67	70	71	71	12
993D2	Armstrong-Gara loams, 9 to 14 percent slopes, moderately eroded	6.39	13.8%		IVe	131.2	38	23	20	62	62	38
1313E3	Munterville soils, 14 to 18 percent slopes, severely eroded	6.99	15.1%		VIIe	80	23.2	15	5	32	32	19
76C2	Ladoga silt loam, dissected till plain, 5 to 9 percent slopes, eroded	7.75	16.7%		llle	192	55.7	75	65	64	64	47

\*\*IA has updated the CSR values for each county to CSR2.

\*- CSR weighted average cannot be calculated on the current soils data, use prior data version for csr values.

\*i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University.

\*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.