



# Soils Map



State: **Missouri**  
 County: **Chariton**  
 Location: **16-53N-19W**  
 Township: **Brunswick**  
 Acres: **145.8**  
 Date: **11/3/2020**

**ENYEART**  
 Auction & Realty  
 Daniel W. Enyeart, Broker/Owner, Auctioneer, MO General Certified Appraiser, 655-451-1161, 655-209-3022  
 112 East Jackson - Hwy 24, Kirksville, MO 65261, www.eneartauctionrealty.com  
 Loren D. Enyeart, Real Estate Salesperson, Real Estate Appraiser, 655-451-9899

Maps Provided By:  
**surety**  
 CUSTOMIZED ONLINE MAPPING  
 © AgriData, Inc. 2020 www.AgriDataInc.com



| Area Symbol: MO041, Soil Area Version: 23 |                                                                                   |       |                  |                      |                         |                  |            |            |              |                  |
|-------------------------------------------|-----------------------------------------------------------------------------------|-------|------------------|----------------------|-------------------------|------------------|------------|------------|--------------|------------------|
| Code                                      | Soil Description                                                                  | Acres | Percent of field | NCCPI Overall Legend | Soil Drainage           | Non-Irr Class *c | Corn       | Soybeans   | Winter wheat | *n NCCPI Overall |
| 64029                                     | Tina silt loam, 0 to 2 percent slopes, rarely flooded                             | 75.55 | 51.8%            |                      | Somewhat poorly drained | IIw              |            |            |              | 83               |
| 36061                                     | Carlow silty clay, 0 to 2 percent slopes, rarely flooded                          | 61.75 | 42.4%            |                      | Poorly drained          | IIIw             |            |            |              | 56               |
| 66111                                     | Waldron silty clay, loamy substratum, 0 to 2 percent slopes, occasionally flooded | 8.34  | 5.7%             |                      | Somewhat poorly drained | IIIw             | 5          | 2          | 2            | 47               |
| 13598                                     | Booker silty clay, frequently ponded, 0 to 2 percent slopes, occasionally flooded | 0.09  | 0.1%             |                      | Very poorly drained     | Vw               |            |            |              | 23               |
| 99001                                     | Water                                                                             | 0.07  | 0.0%             |                      |                         | VIII             |            |            |              |                  |
| <b>Weighted Average</b>                   |                                                                                   |       |                  |                      |                         |                  | <b>0.3</b> | <b>0.1</b> | <b>0.1</b>   | <b>*n 69.4</b>   |

\*n: The aggregation method is "Weighted Average using all components"

\*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.