



# FSA Tillable Soils Map



Soils data provided by USDA and NRCS.



State: **Iowa**  
 County: **Dallas**  
 Location: **3-81N-27W**  
 Township: **Beaver**  
 Acres: **65.01**  
 Date: **9/29/2021**



Maps Provided By:



Area Symbol: IA049, Soil Area Version: 26

Code	Soil Description	Acres	Percent of field	CSR2 Legend	Non-Irr Class *c	*i Corn	*i Soybeans	CSR2**	CSR	*n NCCPI Overall
823B	Ridgeport sandy loam, 2 to 5 percent slopes	18.68	28.7%		IIIe	80	23.2	51	35	56
L138B	Clarion loam, Bemis moraine, 2 to 6 percent slopes	10.22	15.7%		Ile	80	23.2	88		80
L138C2	Clarion loam, Bemis moraine, 6 to 10 percent slopes, moderately eroded	6.90	10.6%		IIIe	0	0	83		67
203	Cylinder loam, 0 to 2 percent slopes	6.51	10.0%		IIs	80	23.2	58	80	73
L107	Webster clay loam, Bemis moraine, 0 to 2 percent slopes	5.71	8.8%		IIw	0	0	88		84
27B	Terril loam, 2 to 6 percent slopes	4.84	7.4%		Ile	224	65	87	82	88
135	Coland clay loam, 0 to 2 percent slopes, occasionally flooded	3.88	6.0%		IIw	198.4	57.5	76	80	83
259	Biscay clay loam, 0 to 2 percent slopes	3.82	5.9%		IIw	166.4	48.3	52	77	69
828C2	Zenor sandy loam, 5 to 9 percent slopes, moderately eroded	2.72	4.2%		IIIe	80	23.2	45	31	49
L55	Nicollet loam, 1 to 3 percent slopes	1.73	2.7%		Ie	0	0	91		85
<b>Weighted Average</b>						<b>85.2</b>	<b>24.7</b>	<b>69.2</b>	<b>*-</b>	<b>*n 70.3</b>

\*\*IA has updated the CSR values for each county to CSR2.

\*- CSR weighted average cannot be calculated on the current soils data, use prior data version for csr values.

\*i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University.

\*n: The aggregation method is "Weighted Average using all components"

\*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.