Tillable Soils Map



	Weighted Average					159.7	46.3	72.4	80.3	*n 80.2
1314	Hanlon-Spillville complex, channeled, 0 to 2 percent slopes	0.08	0.2%		Vw	120.0	34.8	33	25	48
308	Wadena loam, 0 to 2 percent slopes	0.72	1.4%		lls	80.0	23.2	56	73	62
259	Biscay clay loam, 0 to 2 percent slopes	5.30	10.6%		llw	166.4	48.3	52	77	69
135	Coland clay loam, 0 to 2 percent slopes, occasionally flooded	6.99	14.0%		llw	198.4	57.5	76	80	83
536	Hanlon fine sandy loam, 0 to 2 percent slopes	12.01	24.1%		lls	171.2	49.6	79	72	86
203	Cylinder loam, 0 to 2 percent slopes	12.30	24.7%		lls	80.0	23.2	58	80	72
	Spillville loam, 0 to 2 percent slopes, occasionally flooded	12.40	24.9%		llw	208.0	60.3	88	91	87

**IA has updated the CSR values for each county to CSR2.

*i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University.

*n: The aggregation method is "Weighted Average using all components"

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.