

TILLABLE SOILS



use and waives any and all claims against FarmWorth
LLC for any inaccuracies or inconsistencies in the
information and/or data.

Code	Description	Acres	% of field	IA CSR2	Overall NCCPI	Soil Class (Irr)	Soil Class (Non-Irr)	Drainage Class
507	Canisteo clay loam, 0 to 2 percent slopes	31.88	43.64 %	84	82	0	2	Poorly drained
138C2 📘	Clarion loam, 6 to 10 percent slopes, moderately eroded	19.98	27.35 %	83	68	0	3	Well drained
62D2 📃	Storden loam, 10 to 16 percent slopes, moderately eroded	14.05	19.23 %	41	64	0	4	Well drained
107	Webster clay loam, 0 to 2 percent slopes	4.68	6.41 %	86	83	0	2	Poorly drained
956 🗌	Harps-Okoboji complex, 0 to 2 percent slopes	4.31	5.90 %	69	79	0	2	Poorly drained
27B	Terril loam, 2 to 6 percent slopes	2.00	2.74 %	87	91	0	2	Well drained
62C2 🗌	Storden loam, 6 to 10 percent slopes, moderately eroded	0.45	0.62 %	64	67	0	3	Well drained
638C2 📘	Clarion-Storden complex, 6 to 10 percent slopes, moderately eroded	0.02	0.03 %	75	68	0	3	Well drained
138B 📃	Clarion loam, 2 to 6 percent slopes	0.01	0.01 %	89	83	0	2	Moderately well drained
Average:					75.2			

IA has updated the CSR values for each county to CSR2. Soils data provided by USDA and NRCS.