

## RIVERVIEW FAMILY FARMS - BLACK HAWK (289 AC TOTAL, 280.05 AC TILLABLE)

## TILLABLE SOILS





Field borders obtained from Farm Service Agency as of 2008

FarmWorth, LLC makes no representations or warranties, express or implied, as to the accuracy of any information, data, numerical values, boundaries, or any other information generated through the use of FarmWorth.com. User is solely responsible for independently investigating and determining all information provided through FarmWorth.com prior to use and waives any and all claims against FarmWorth, LLC for any inaccuracies or inconsistencies in the information and/or data.

Code	Description	Acres	% of field	IA CSR2	Soil Class (Non-Irr)	Drainage Class
84	Clyde silty clay loam, 0 to 3 percent slopes	88.25	31.51 %	88	2	Poorly drained
468B 📃	Dunkerton sandy loam, 2 to 5 percent slopes	63.62	22.72 %	76	2	Somewhat poorly drained
471	Oran loam, 1 to 3 percent slopes	35.01	12.50 %	74	1	Somewhat poorly drained
63B 📕	Chelsea loamy fine sand, 2 to 5 percent slopes	29.13	10.40 %	19	4	Excessively drained
391B 📃	Clyde-Floyd complex, 1 to 4 percent slopes	21.76	7.77 %	87	2	Poorly drained
171B 📃	Bassett loam, 2 to 5 percent slopes	11.39	4.07 %	85	2	Moderately well drained
198B 📃	Floyd loam, 1 to 4 percent slopes	11.21	4.00 %	89	2	Somewhat poorly drained
41B 📃	Sparta loamy sand, 2 to 5 percent slopes	8.98	3.21 %	42	4	Excessively drained
41	Sparta loamy sand, 0 to 2 percent slopes	5.54	1.98 %	48	4	Excessively drained
63C 📕	Chelsea loamy fine sand, 5 to 9 percent slopes	3.03	1.08 %	14	4	Excessively drained
408B 📃	Olin sandy loam, 2 to 5 percent slopes	2.10	0.75 %	64	2	Well drained
585	Coland-Spillville complex, 0 to 2 percent slopes, occasionally flooded	0.01	0.00 %	70	2	Poorly drained
			Average:	72.9		

IA has updated the CSR values for each county to CSR2.

Soils data provided by USDA and NRCS.