Soils Map



	Symbol: IA083, Soil Area Version		1				1	1				1
Code	Soil Description	Acres	Percent of field	CSR2 Legend	Non-Irr Class *c	*i Corn Bu	*i Soybeans Bu	CSR2**	CSR	Oats Bu	Spring wheat Bu	*n NCCPI Soybeans
177	Saude loam, 0 to 2 percent slopes	20.02	33.4%		lls	80.0	23.2	60	63			62
485	Spillville loam, 0 to 2 percent slopes, occasionally flooded	19.41	32.3%		llw	208.0	60.3	88	92			87
135	Coland clay loam, 0 to 2 percent slopes, occasionally flooded	5.06	8.4%		llw	198.4	57.5	76	80			83
828B	Zenor sandy loam, 1 to 5 percent slopes	4.15	6.9%		llle	80.0	23.2	51	51			32
936	Coland-Spillville-Hanlon complex, 0 to 2 percent slopes	3.97	6.6%		llw	200.0	58.0	82	75			87
73E	Hawick gravelly sandy loam, 12 to 20 percent slopes	2.51	4.2%		VIIs	80.0	23.2	5	5			18
1226	Lawler loam, 0 to 2 percent slopes, rarely flooded	2.46	4.1%		lls	80.0	23.2	59	73			60
27B	Terril loam, 2 to 6 percent slopes	1.66	2.8%		lle	224.0	65.0	87	84			84
236F	Lester loam, Bemis moraine, 10 to 22 percent slopes	0.40	0.7%		Vle	129.6	37.6	36	23	2	1	58
201B	Coland-Terril complex, 2 to 5 percent slopes	0.23	0.4%		llw	208.0	60.3	80	69			90



Code	Soil Description	Acres	Percent of field	-		-	*i Soybeans Bu	CSR2**	CSR		Spring wheat Bu	*n NCCPI Soybeans
178B	Waukee loam, 2 to 5 percent slopes	0.15	0.2%		lls	80.0	23.2	64	74			69
Weighted Average					2.30	144.1	41.8	69.6	72.1	*-	*-	*n 70.2

**IA has updated the CSR values for each county to CSR2. *i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University. *n: The aggregation method is "Weighted Average using all components" *c: Using Capabilities Class Dominant Condition Aggregation Method