





153AC, CHICKASAW CO  
TOTAL SOILS



County: Chickasaw County, IA  
Location: 24-94N-14W  
Township: Bradford  
Acres: 152.64  
Date: 09/05/2024



Field borders obtained from Farm Service Agency  
as of 2008

FarmWorth, LLC makes no representations or warranties, express or implied, as to the accuracy of any information, data, numerical values, boundaries, or any other information generated through the use of FarmWorth.com. User is solely responsible for independently investigating and determining all information provided through FarmWorth.com prior to use and waives any and all claims against FarmWorth, LLC for any inaccuracies or inconsistencies in the information and/or data.

| Code     | Description                                                            | Acres | % of field | IA CSR2 | Soil Class (Non-Irr) | Drainage Class               |
|----------|------------------------------------------------------------------------|-------|------------|---------|----------------------|------------------------------|
| 391B     | Clyde-Floyd complex, 1 to 4 percent slopes                             | 41.85 | 27.42 %    | 87      | 2                    | Poorly drained               |
| 83B      | Kenyon loam, 2 to 5 percent slopes                                     | 38.91 | 25.49 %    | 90      | 2                    | Moderately well drained      |
| 399      | Readlyn silt loam, 1 to 3 percent slopes                               | 34.20 | 22.41 %    | 91      | 1                    | Somewhat poorly drained      |
| 84       | Clyde clay loam, 0 to 3 percent slopes                                 | 10.97 | 7.19 %     | 88      | 2                    | Poorly drained               |
| 783C     | Cresco loam, 5 to 9 percent slopes                                     | 8.31  | 5.44 %     | 81      | 3                    | Moderately well drained      |
| 83C2     | Kenyon loam, 5 to 9 percent slopes, eroded                             | 6.73  | 4.41 %     | 84      | 3                    | Moderately well drained      |
| 198B     | Floyd loam, 1 to 4 percent slopes                                      | 3.86  | 2.53 %     | 89      | 2                    | Somewhat poorly drained      |
| 471B     | Oran loam, 2 to 5 percent slopes                                       | 3.08  | 2.02 %     | 74      | 1                    | Somewhat poorly drained      |
| 284B     | Flagler sandy loam, 2 to 5 percent slopes                              | 2.16  | 1.42 %     | 49      | 3                    | Somewhat excessively drained |
| 585      | Coland-Spillville complex, 0 to 2 percent slopes, occasionally flooded | 1.71  | 1.12 %     | 70      | 2                    | Poorly drained               |
| 783B     | Cresco loam, 2 to 5 percent slopes                                     | 0.57  | 0.37 %     | 73      | 2                    | Moderately well drained      |
| 798B     | Protivin loam, 1 to 4 percent slopes                                   | 0.31  | 0.20 %     | 61      | 2                    | Somewhat poorly drained      |
| Average: |                                                                        |       |            | 87.2    |                      |                              |

IA has updated the CSR values for each county to CSR2.  
Soils data provided by USDA and NRCS.