



PLUM CREEK TWP (80.00 AC TOTAL, 79.79 AC TILLABLE)

TILLABLE SOILS





County: Kossuth County, IA
Location: 34-96N-28W
Township: Plum Creek

Acres: 79.79
Date: 10/04/2024



Field borders obtained from Farm Service Agency as of 2008

FarmWorth, LLC makes no representations or warranties, express or implied, as to the accuracy of any information, data, numerical values, boundaries, or any other information generated through the use of FarmWorth.com. User is solely responsible for independently investigating and determining all information provided through FarmWorth.com prior to use and waives any and all claims against FarmWorth, LLC for any inaccuracies or inconsistencies in the information and/or data.

| Code | Description | Acres | % of field | IA CSR2 | Overall NCCPI | Soil Class (Irr) | Soil Class (Non- Irr) | Drainage Class |
|------|--|-------|---------------|------------|------------------|---------------------|--------------------------|----------------------------|
| 221 | Klossner muck, 0 to 1 percent slopes | 38.43 | 48.16 % | 32 | 84 | - | 3 | Very poorly drained |
| 95 | Harps clay loam, 0 to 2 percent slopes | 10.05 | 12.60 % | 72 | 82 | - | 2 | Poorly drained |
| 90 | Okoboji mucky silt loam, 0 to 1 percent slopes | 10.02 | 12.56 % | 56 | 70 | - | 3 | Very poorly drained |
| 55 | Nicollet clay loam, 1 to 3 percent slopes | 9.03 | 11.32 % | 89 | 81 | - | 1 | Somewhat poorly drained |
| 507 | Canisteo clay loam, 0 to 2 percent slopes | 8.11 | 10.16 % | 84 | 82 | - | 2 | Poorly drained |
| 6 🗌 | Okoboji silty clay loam, 0 to 1 percent slopes | 2.38 | 2.98 % | 59 | 75 | - | 3 | Very poorly drained |
| 138B | Clarion loam, 2 to 6 percent slopes | 1.01 | 1.27 % | 89 | 83 | - | 2 | Moderately well drained |
| 107 | Webster clay loam, 0 to 2 percent slopes | 0.75 | 0.94 % | 86 | 83 | - | 2 | Poorly drained |
| | | | Average: | 53.8 | 81.2 | | | |

IA has updated the CSR values for each county to CSR2. Soils data provided by USDA and NRCS.