







Field borders obtained from Farm Service Agency as of 2008

FarmWorth, LLC makes no representations or warranties, express or implied, as to the accuracy of any information, data, numerical values, boundaries, or any other information generated through the use of FarmWorth.com. User is solely responsible for independently investigating and determining all information provided through FarmWorth.com prior to use and waives any and all claims against FarmWorth, LLC for any inaccuracies or inconsistencies in the information and/or data.

Code	Description	Acres	% of field	IA CSR2	Soil Class (Non-Irr)	Drainage Class
83B 📃	Kenyon loam, 2 to 5 percent slopes	49.86	30.87 %	90	2	Moderately well drained
198B 📃	Floyd loam, 1 to 4 percent slopes	41.90	25.94 %	89	2	Somewhat poorly drained
84	Clyde clay loam, 0 to 3 percent slopes	30.94	19.16 %	88	2	Poorly drained
221 📃	Klossner muck, 1 to 4 percent slopes	8.67	5.37 %	48	3	Very poorly drained
83C2 📃	Kenyon loam, 5 to 9 percent slopes, eroded	8.14	5.04 %	84	3	Moderately well drained
171B 📃	Bassett loam, 2 to 5 percent slopes	4.80	2.97 %	85	2	Moderately well drained
399 📃	Readlyn silt loam, 1 to 3 percent slopes	4.30	2.66 %	91	1	Somewhat poorly drained
C135 📕	Coland clay loam, channeled, 0 to 2 percent slopes	3.10	1.92 %	18	5	Poorly drained
1226 📃	Lawler loam, 0 to 2 percent slopes, rarely flooded	2.64	1.63 %	59	2	Somewhat poorly drained
284 🗌	Flagler sandy loam, 0 to 2 percent slopes	2.59	1.60 %	56	3	Somewhat excessively drained
409B 📃	Dickinson fine sandy loam, till substratum, 2 to 5 percent slopes	2.40	1.49 %	74	3	Well drained
394B 📃	Ostrander loam, 2 to 5 percent slopes	2.17	1.34 %	88	2	Well drained
			Average:	84.0		

IA has updated the CSR values for each county to CSR2. Soils data provided by USDA and NRCS.