

LOFSHULT (162.98 AC TOTAL, 163.82 AC TILLABLE)

TILLABLE SOILS





Field borders obtained from Farm Service Agency as of 2008

FarmWorth, LLC makes no representations or warranties, express or implied, as to the accuracy of any information, data, numerical values, boundaries, or any other information generated through the use of FarmWorth.com. User is solely responsible for independently investigating and determining all information provided through FarmWorth.com prior to use and waives any and all claims against FarmWorth, LLC for any inaccuracies or inconsistencies in the information and/or data.

Code	Description	Acres	% of field	IA CSR2	Overall NCCPI	Soil Class (Irr)	Soil Class (Non- Irr)	Drainage Class
Couc	Beschpilon	7 101 00	noid	COIL		(,	,	Dramage class
67 📘	Woodbury silty clay, 0 to 2 percent slopes, rarely flooded	102.21	62.39 %	70	53	-	3	Poorly drained
436 📃	Lakeport silty clay loam, 0 to 2 percent slopes, rarely flooded	44.65	27.26 %	88	70	-	1	Somewhat poorly drained
66 🗌	Luton silty clay, 0 to 2 percent slopes, rarely flooded	16.96	10.35 %	57	51	-	3	Poorly drained
Average:					57.4			

IA has updated the CSR values for each county to CSR2.

Soils data provided by USDA and NRCS.