## Soils Map



Code	Soil Description	Acres	Percent of field	CSR2 Legend	Non-Irr Class *c	*i Corn Bu	*i Soybeans Bu	CSR2**	CSR	*n NCCPI Soybeans
L329	Webster-Nicollet complex, Bemis moraine, 0 to 3 percent slopes	116.66	36.4%		llw	0.0	0.0	89		83
L95	Harps clay loam, Bemis moraine, 0 to 2 percent slopes	97.95	30.6%		llw	0.0	0.0	75		80
L138B	Clarion loam, Bemis moraine, 2 to 6 percent slopes	38.96	12.2%		lle	220.8	64.0	88		79
6	Okoboji silty clay loam, 0 to 1 percent slopes	30.60	9.5%		IIIw	185.6	53.8	59	59	74
L507	Canisteo clay loam, Bemis moraine, 0 to 2 percent slopes	12.94	4.0%		llw	0.0	0.0	87		81
L107	Webster clay loam, Bemis moraine, 0 to 2 percent slopes	9.21	2.9%		llw	0.0	0.0	88		83
1226	Lawler loam, 0 to 2 percent slopes, rarely flooded	8.17	2.5%		lls	80.0	23.2	59	73	60
L55	Nicollet loam, 1 to 3 percent slopes	6.05	1.9%		le	0.0	0.0	91		84
Weighted Average						46.6	13.5	80.9	*-	*n 80.1

\*\*IA has updated the CSR values for each county to CSR2.

\*n: The aggregation method is "Weighted Average using all components"

\*c: Using Capabilities Class Dominant Condition Aggregation Method

<sup>\*-</sup> CSR weighted average cannot be calculated on the current soils data, use prior data version for csr values.\*i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University.