## Soils Map



Area Syn	nbol: IA083, Soil Area Version: 31									
Code	Soil Description	Acres	Percent of field	CSR2 Legend	Non-Irr Class *c	*i Corn Bu	*i Soybeans Bu	CSR2**	CSR	*n NCCPI Soybeans
L329	Webster-Nicollet complex, Bemis moraine, 0 to 3 percent slopes	61.68	38.5%		llw	0.0	0.0	89		83
L95	Harps clay loam, Bemis moraine, 0 to 2 percent slopes	39.61	24.7%		llw	0.0	0.0	75		80
L138B	Clarion loam, Bemis moraine, 2 to 6 percent slopes	17.09	10.7%		lle	220.8	64.0	88		79
L638C2	Clarion-Storden complex, Bemis moraine, 6 to 10 percent slopes, moderately eroded	8.94	5.6%		llle	0.0	0.0	75		67
L55	Nicollet loam, 1 to 3 percent slopes	8.88	5.5%		le	0.0	0.0	91		84
L507	Canisteo clay loam, Bemis moraine, 0 to 2 percent slopes	7.35	4.6%		llw	0.0	0.0	87		81
177B	Saude loam, 2 to 5 percent slopes	6.06	3.8%		lls	80.0	23.2	55	58	60
L138C2	Clarion loam, Bemis moraine, 6 to 10 percent slopes, moderately eroded	4.68	2.9%		llle	0.0	0.0	83		64
6	Okoboji silty clay loam, 0 to 1 percent slopes	3.74	2.3%		IIIw	185.6	53.8	59	59	74
27B	Terril loam, 2 to 6 percent slopes	2.16	1.3%		lle	224.0	65.0	87	84	84
L107	Webster clay loam, Bemis moraine, 0 to 2 percent slopes	0.21	0.1%		llw	0.0	0.0	88		83
Weighted Average						33.9	9.8	82.5	*-	*n 79.3

\*\*IA has updated the CSR values for each county to CSR2.

\*- CSR weighted average cannot be calculated on the current soils data, use prior data version for csr values.\*i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University.

\*n: The aggregation method is "Weighted Average using all components"

\*c: Using Capabilities Class Dominant Condition Aggregation Method