



County: Palo Alto County, IA
Location: 22-95N-34W

Township: Silver Lake
Acres: 91.62

Date: **04/15/2025**



Field borders obtained from Farm Service Agency as of 2008

FarmWorth, LLC makes no representations or warranties, express or implied, as to the accuracy of any information, data, numerical values, boundaries, or any other information generated through the use of FarmWorth.com. User is solely responsible for independently investigating and determining all information provided through FarmWorth.com prior to use and waives any and all claims against FarmWorth, LLC for any inaccuracies or inconsistencies in the information and/or data.

Code	Description	Acres	% of field	IA CSR2	Overall NCCPI	Soil Class (Irr)	Soil Class (Non-Irr)	Drainage Class
138B	Clarion loam, 2 to 6 percent slopes	28.77	31.40 %	89	83	-	2	Moderately well drained
55	Nicollet clay loam, 1 to 3 percent slopes	19.29	21.06 %	89	81	-	1	Somewhat poorly drained
133	Colo silty clay loam, 0 to 2 percent slopes	14.71	16.06 %	74	82	-	2	Poorly drained
138C2	Clarion loam, 6 to 10 percent slopes, moderately eroded	9.74	10.63 %	83	68	-	3	Well drained
339B	Truman silt loam, 2 to 6 percent slopes	7.59	8.28 %	95	89	-	2	Well drained
73C	Hawick gravelly sandy loam, 6 to 12 percent slopes	5.87	6.40 %	31	37	-	4	Excessively drained
62D 🔲	Storden loam, 10 to 16 percent slopes, moderately eroded	3.48	3.80 %	41	64	-	4	Well drained
133B	Colo silty clay loam, 2 to 4 percent slopes	1.04	1.13 %	80	97	-	2	Poorly drained
6 🔲	Okoboji silty clay loam, 0 to 1 percent slopes	0.67	0.73 %	59	75	-	3	Very poorly drained
181C2 🔲	Clarion-Estherville complex, 5 to 9 percent slopes, moderately eroded	0.35	0.38 %	49	55	-	4	Well drained
507	Canisteo clay loam, 0 to 2 percent slopes	0.12	0.13 %	84	82	-	2	Poorly drained
			Average:	80.4	77.6			

IA has updated the CSR values for each county to CSR2. Soils data provided by USDA and NRCS.