



## EWING FARM (30.75 AC TOTAL, 17.69 AC TILLABLE)

## **TOTAL SOILS**





County: Wayne County, IA Location: 09-70N-20W

Township: Wright Acres: 30.75

Date: 05/01/2025



Field borders obtained from Farm Service Agency as of 2008

FarmWorth, LLC makes no representations or warranties, express or implied, as to the accuracy of any information, data, numerical values, boundaries, or any other information generated through the use of FarmWorth.com. User is solely responsible for independently investigating and determining all information provided through FarmWorth.com prior to use and waives any and all claims against FarmWorth, LLC for any inaccuracies or inconsistencies in the information and/or data.

| Code    | Description  | Acres | % of<br>field | IA<br>CSR2 | Overall<br>NCCPI | Soil Class<br>(Irr) | Soil Class<br>(Non-Irr) | Drainage Class          |
|---------|--|-------|---------------|------------|------------------|---------------------|-------------------------|-------------------------|
| 179E2   | Gara loam, 14 to 18 percent slopes, moderately eroded                    | 9.32  | 30.30 %       | 24         | 62               | -                   | 6                       | Well drained            |
| 711     | Lawson-Nodaway-Colo complex, 0 to 2 percent slopes, occasionally flooded | 5.22  | 16.97 %       | 83         | 88               | -                   | 2                       | Somewhat poorly drained |
| 792C2   | Armstrong loam, 5 to 9 percent slopes, moderately eroded                 | 4.87  | 15.83 %       | 31         | 66               | -                   | 3                       | Somewhat poorly drained |
| 269B 🔲  | Humeston silty clay loam, 2 to 5 percent slopes, rarely flooded          | 4.23  | 13.75 %       | 71         | 89               | -                   | 3                       | Poorly drained          |
| 179D2 📕 | Gara loam, 9 to 14 percent slopes, moderately eroded                     | 2.73  | 8.88 %        | 38         | 66               | -                   | 4                       | Well drained            |
| 13B 🔲   | Zook-Olmitz-Vesser complex, 0 to 5 percent slopes                        | 2.50  | 8.13 %        | 68         | 77               | -                   | 2                       | Poorly drained          |
| 531C 🗌  | Kniffin silt loam, 5 to 9 percent slopes                                 | 0.88  | 2.86 %        | 52         | 67               | -                   | 3                       | Somewhat poorly drained |
| 531C2 🔲 | Kniffin silty clay loam, 5 to 9 percent slopes, moderately eroded        | 0.55  | 1.79 %        | 48         | 61               | -                   | 3                       | Somewhat poorly drained |
| 312B 🔲  | Seymour silt loam, 2 to 5 percent slopes                                 | 0.46  | 1.50 %        | 64         | 70               | -                   | 3                       | Somewhat poorly drained |
|         | Average:   |       |               |            | 72.6             |                     |                         |                         |

IA has updated the CSR values for each county to CSR2. Soils data provided by USDA and NRCS.